arXiv:1101.4931v1 [math.NT] 24 Jan 2011

On q-Bernstein and q-Hermite polynomials

T. Kim[1], J. Choi[1], Y. H. Kim[1], C. S. Ryoo[2]

[1]Division of General Education-Mathematics,
Kwangwoon University, Seoul 139-701, Korea

[2]Department of Mathematics,
Hannam University, Daejeon 306-791, Korea

Abstract Recently, T. Kim([1,2]) introduced q-Bernstein polynomials which are different q-Bernstein polynomials of Phillips. In this paper, we investigate some identities for the q-Bernstein polynomials which are treated by Kim in previous paper([1]).

2000 Mathematics Subject Classification: 11B68, 11S40, 11S80

Key words- Bernstein polynomials, Hermite polynomials, q-Bernstein polynomials, q-Hermite polynomials

1. Introduction

Let $C[0,1]$ denote the set of continuous functions on $[0,1]$. For $f \in C[0,1]$, Bernstein introduced the following well-known linear operator(see [3, 5, 8]):

$$\begin{aligned}\mathbb{B}_n(f \mid x) &= \sum_{k=0}^{n} f\left(\frac{k}{n}\right)\binom{n}{k}x^k(1-x)^{n-k} \\ &= \sum_{k=0}^{n} f\left(\frac{k}{n}\right)B_{k,n}(x),\end{aligned} \tag{1}$$

where $\binom{n}{k} = \dfrac{n(n-1)\cdots(n-k+1)}{n!}$. Here $\mathbb{B}_n(f \mid x)$ is called the Bernstein operator of order n for f. For $n, k \in \mathbb{Z}_+ = \mathbb{N} \cup \{0\}$, the Bernstein polynomial of degree n is defined by

$$B_{k,n}(x) = \binom{n}{k}x^k(1-x)^{n-k}, \text{ (see [1-8])} .$$

The Hermite polynomials have the generating function

$$e^{2zw-w^2}=\sum_{n=0}^{\infty}\frac{H_n(z)}{n!}w^n,\quad z,w\in\mathbb{C},\tag{2}$$

which gives the Cauchy type integral

$$H_n(z)=\frac{n!}{2\pi i}\int_C e^{2zw-w^2}\frac{d^w}{w^{n+1}},$$

where C is a circle around the origin with the positive direction (see [4]).

From (2), we note that the Hermite polynomials are given by

$$H_n(x)=n!\sum_{k=0}^{[\frac{n}{2}]}\frac{(-1)^k}{k!(n-2k)!}(2x)^{n-2k},$$

where $[\,\cdot\,]$ is the Gauss symbol (see [4]).

Let $0<q<1$. Define the q-number of x by

$$[x]_q=\frac{1-q^x}{1-q}.$$

Note that $\lim_{q\to 1}[x]_q=x$ (see [9-11]).

For $f\in C[0,1]$, the q-Bernstein type operator of order n for f is defined by

$$\begin{aligned}\mathbb{B}_{n,q}(f\mid x)&=\sum_{k=0}^{n}f\left(\frac{k}{n}\right)\binom{n}{k}[x]_q^k[1-x]_{\frac{1}{q}}^{n-k}\\&=\sum_{k=0}^{n}\left(\frac{k}{n}\right)B_{k,n}(x\mid q),\end{aligned}\tag{4}$$

where $n,k\in\mathbb{Z}_+$ (see [1]). Here $B_{k,n}(x\mid q)=\binom{n}{k}[x]_q^k[1-x]_{\frac{1}{q}}^{n-k}$ are called the q-Bernstein type polynomials of degree n (see [1, 2]). Note that (4) is a q-analogue of (1).

In this paper, we consider the q-Hermite polynomials related to the q-Bernstein type polynomials. From these q-Hermite polynomials, we give some interesting identities for the q-Bernstein polynomials.

2. q-Bernstein polynomials and q-Hermite polynomials

In this section, we consider a q-analogue of (2), which are called the q-Hermite polynomials, as follows:

$$e^{2t[x]_q-t^2}=\sum_{n=0}^{\infty}H_{n,q}(x)\frac{t^n}{n!}.\tag{5}$$

From (5), we have

$$\begin{aligned}
e^{2t[x]_q-t^2} &= \left(\sum_{m=0}^{\infty}\frac{(2[x]_q)^m}{m!}t^m\right)\left(\sum_{l=0}^{\infty}\frac{(-1)^l}{l!}t^{2l}\right) \\
&= \sum_{n=0}^{\infty}\left(\sum_{l=0}^{\left[\frac{n}{2}\right]}\frac{(-1)^l(2[x]_q)^{n-2l}n!}{l!(n-2l)!}\right)\frac{t^n}{n!}.
\end{aligned} \tag{6}$$

Therefore, by (5) and (6), we obtain the following theorem.

Theorem 1. For $n \in \mathbb{Z}_+$, we have

$$H_{n,q}(x) = n!\sum_{l=0}^{\left[\frac{n}{2}\right]}\frac{(-1)^l 2^{n-2l}[x]_q^{n-2l}}{l!(n-2l)!}.$$

Note that Theorem 1 is a q-analogue of (3). By (4), we obtain the following corollary.

Corollary 2. For $x \in [0,1]$, we have

$$H_{n,q}(x) = n!\sum_{l=0}^{\left[\frac{n}{2}\right]}\frac{(-1)^l 2^{n-2l}B_{2l,n}(1-x \mid \frac{1}{q})}{l!(n-2l)![1-x]_{\frac{1}{q}}^{2l}}.$$

From the definition of q-Bernstein polynomials, we note that

$$\begin{aligned}
\sum_{n=0}^{\infty}\frac{(-1)^n}{n!}\left(\frac{d}{d[x]_q}\right)^n e^{-t[x]_q} &= \sum_{n=0}^{\infty}\frac{(-1)^n}{n!}(-t)^n e^{-t[x]_q} \\
&= \left(\sum_{n=0}^{\infty}\frac{t^n}{n!}\right)e^{-t[x]_q} = e^{[1-x]_{\frac{1}{q}}t} \\
&= \frac{k!}{(t[x]_q)^k}\left(\frac{(t[x]_q)^k e^{[1-x]_{\frac{1}{q}}t}}{k!}\right) \\
&= \frac{k!}{(t[x]_q)^k}\sum_{n=k}^{\infty}B_{k,n}(x \mid q)\frac{t^n}{n!} \\
&= \frac{k!}{(t[x]_q)^k}\sum_{n=0}^{\infty}B_{k,n+k}(x \mid q)\frac{t^{n+k}}{(n+k)!} \\
&= \frac{1}{[x]_q^k}\sum_{n=0}^{\infty}\frac{k!}{(n+k)\cdots(n+1)}B_{k,n+k}(x \mid q)\frac{t^n}{n!} \\
&= \frac{1}{[x]_q^k}\sum_{n=0}^{\infty}\frac{B_{k,n+k}(x \mid q)}{\binom{n+k}{k}}\frac{t^n}{n!}.
\end{aligned} \tag{7}$$

By a simple calculation, we easily get

$$\begin{aligned}\sum_{n=0}^{\infty}\frac{(-1)^n}{n!}\left(\frac{d}{d[x]_q}\right)^n e^{-t[x]_q} &= \sum_{n=0}^{\infty}\left(\sum_{l=0}^{n}\frac{(-1)^l n!}{l!(n-l)!}\left(\frac{d}{d[x]_q}\right)^l [x]_q^{n-l}\right)\frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty}\left(\sum_{l=0}^{n}\binom{n}{l}(-1)^l\left(\frac{d}{d[x]_q}\right)^l [x]_q^{n-l}\right)\frac{t^n}{n!}.\end{aligned} \tag{8}$$

Therefore, by (7) and (8), we obtain the following theorem.

Theorem 3. For $x \in [0,1], n, k \in \mathbb{Z}_+$, we have

$$[x]_q^k \sum_{l=0}^{n}\binom{n}{l}(-1)^l\left(\frac{d}{d[x]_q}\right)^l [x]_q^{n-l} = \frac{B_{k,n+k}(x \mid q)}{\binom{n+k}{k}}.$$

By a simple calculation, we easily get

$$\left(\frac{1}{2}\frac{d}{d[x]_q}\right)^n e^{-2t[x]_q} = (-t)^n e^{-2t[x]_q}.$$

Thus, we have

$$\sum_{n=0}^{\infty}\frac{(-1)^n}{n!}\left(\frac{1}{2}\frac{d}{d[x]_q}\right)^{2n} e^{-2t[x]_q} = e^{-t^{2n}+2t[x]_q}e^{-4t[x]_q}. \tag{9}$$

From (6) and (9), we can derive the following equation:

$$e^{-t^{2n}-2t[x]_q} = \sum_{n=0}^{\infty}\left(n!(-1)^n\sum_{l=0}^{\left[\frac{n}{2}\right]}\frac{(-1)^l 4^{n-l}[x]_q^{n-l}}{l!(n-2l)!}\right)\frac{t^n}{n!} \tag{10}$$

and

$$e^{-t^2+2t[x]_q}e^{-4t[x]_q} = \sum_{n=0}^{\infty}\left(\sum_{l=0}^{n}\binom{n}{l}H_{l,q}(x)(-1)^{n-l}[x]_q^{n-l}\right)\frac{t^n}{n!}. \tag{11}$$

Therefore, by (10) and (11), we obtain the following theorem.

Theorem 4. For $n \in \mathbb{Z}_+$, we have

$$H_{n,q}(x) = \sum_{l=0}^{n}\binom{n}{l}H_{l,q}(x)(-1)^{n-l}[x]_q^{n-l}.$$

By (4) and Theorem 4, we obtain the following corollary.

Corollary 5. For $n \in \mathbb{Z}_+$ and $x \in [0,1]$, we have

$$H_{n,q}(x) = \sum_{l=0}^{n} \binom{n}{l} H_{l,q}(x)(-1)^l B_{l,n}(1-x \mid \frac{1}{q})[1-x]_{\frac{1}{q}}^{-l}.$$

For $t \in \mathbb{R}$ and $x \in [0,1]$, we have

$$\begin{aligned}
\sinh(t[1-x]_{\frac{1}{q}}) &= \frac{e^{t[1-x]_{\frac{1}{q}}} - e^{-t[1-x]_{\frac{1}{q}}}}{2} \\
&= \frac{k!}{2(t[x]_q)^k}\left(\frac{(t[x]_q)^k e^{t[1-x]_{\frac{1}{q}}} - (t[x]_q)^k e^{-t[1-x]_{\frac{1}{q}}}}{k!}\right) \\
&= \frac{k!}{2t^k[x]_q^k}\left(\sum_{n=k}^{\infty} B_{k,n}(x \mid q)\frac{t^n}{n!} - \sum_{n=k}^{\infty} B_{k,n}(x \mid q)(-1)^{n-k}\frac{t^n}{n!}\right) \\
&= \frac{k!}{2t^k[x]_q^k}\left(\sum_{n=k}^{\infty}(1-(-1)^{n+k})B_{k,n}(x \mid q)\frac{t^n}{n!}\right) \\
&= \frac{k!}{2[x]_q^k}\left(\sum_{n=0}^{\infty}(1-(-1)^{n})B_{k,n+k}(x \mid q)\frac{t^n}{(n+k)!}\right) \\
&= \frac{1}{2[x]_q^k}\left(\sum_{n=0}^{\infty}(1-(-1)^{n})\frac{B_{k,n+k}(x \mid q)}{\binom{n+k}{k}}\frac{t^n}{n!}\right).
\end{aligned} \tag{12}$$

Using Taylor expansion, we get

$$\frac{e^{[1-x]_{\frac{1}{q}}t} - e^{-[1-x]_{\frac{1}{q}}t}}{2} = \sum_{n=0}^{\infty} \frac{[1-x]_{\frac{1}{q}}^n - (-[1-x]_{\frac{1}{q}})^n}{2}\frac{t^n}{n!}. \tag{13}$$

Comparing the coefficients on the both sides of (12) and (13), we obtain the following theorem.

Theorem 6. For $n, k \in \mathbb{Z}_+$ and $x \in [0,1]$, we have

$$[x]_q^{n+k}([1-x]_{\frac{1}{q}}^n - (-1)^n[1-x]_{\frac{1}{q}}^n) = [x]_q^n\big(1-(-1)^n\big)\frac{B_{k,n+k}(x \mid q)}{\binom{n+k}{k}}.$$

From (4) and Theorem 6, we note that

$$\begin{aligned}
&B_{n,2n+k}(1-x \mid \frac{1}{q}) - (-1)^n B_{n,2n+k}(1-x \mid \frac{1}{q}) \\
&= B_{n+k,2n+k}(x \mid q) - (-1)^n B_{n+k,2n+k}(x \mid q).
\end{aligned} \tag{14}$$

Therefore, by (14), we obtain the following theorem.

Theorem 7. For $n, k \in \mathbb{Z}_+$ and $x \in [0,1]$, we have

$$B_{n+k,2n+k}(x \mid q) = [x]_q^n \frac{B_{k,n+k}(x \mid q)}{\binom{n+k}{n}}.$$

By Theorem 7, we get

$$B_{n+1+k,4n+2+k}(x \mid q) = \frac{B_{k,2n+1+k}(x \mid q)}{\binom{2n+1+k}{2n+1}} [x]_q^{2n+1}.$$

Thus, we have

$$\binom{2n+1+k}{2n+1}\binom{4n+2+k}{2n+1+k} = \binom{2n+1+k}{k}.$$

Let $t \in \mathbb{R}$ and $0 \leq x \leq 1$. Then we see that

$$\begin{aligned}
\cosh(t[1-x]_{\frac{1}{q}}) &= \frac{e^{t[1-x]_{\frac{1}{q}}} + e^{-t[1-x]_{\frac{1}{q}}}}{2} \\
&= \frac{k!}{2(t[x]_q)^k} \left(\frac{(t[x]_q)^k}{k!} e^{t[1-x]_{\frac{1}{q}}} + \frac{(-1)^k(-t[x]_q)^k}{k!} e^{-t[1-x]_{\frac{1}{q}}} \right) \\
&= \frac{k!}{2(t[x]_q)^k} \left(\sum_{n=k}^{\infty} B_{k,n}(x \mid q) \frac{t^n}{n!} + \sum_{n=k}^{\infty} B_{k,n}(x \mid q)(-1)^{n-k} \frac{t^n}{n!} \right) \\
&= \frac{1}{2[x]_q^k} \left(\sum_{n=0}^{\infty} (1+(-1)^n) \frac{B_{k,n+k}(x \mid q)}{\binom{n+k}{k}} \frac{t^n}{n!} \right).
\end{aligned} \tag{15}$$

Using Taylor expansion, we get

$$\cosh(t[1-x]_{\frac{1}{q}}) = \frac{1}{2} \sum_{n=0}^{\infty} (1+(-1)^n)[1-x]_{\frac{1}{q}}^n \frac{t^n}{n!}. \tag{16}$$

Therefore, by (15) and (16), we obtain

$$[x]_q^{n+k}[1-x]_{\frac{1}{q}}^n = [x]_q^n \frac{B_{k,n+k}(x \mid q)}{\binom{n+k}{k}}.$$

Thus, we have

$$B_{2n+k,4n+k}(x \mid q) = \frac{B_{k,2n+k}(x \mid q)}{\binom{2n+k}{2n}} [x]_q^{2n}. \tag{17}$$

From (17), we have

$$\binom{4n+k}{2n+k}\binom{2n+k}{2n} = \binom{2n+k}{k}.$$

References

[1] T. Kim, A note on q-Bernstein polynomials, Russian Journal of Mathematical physics, 18(2011), no.1.

[2] T. Kim, J. Choi, Y. H. Kim, q-Bernstein polynomials associated with q-Stirling numbers and Carlitz's q-Bernoulli numbers, Abstract and Applied Analysis , 2010(2010), Article ID 150975, 11 pages.

[3] T. Kim, L.-C. Jang, H. Yi, A note on the modified q-Bernstein polynomials, Discrete Dynamics in Nature and Society, 2010(2010), Article ID 706483, 12 pages.

[4] J. L. Lopez, N. M. Temme, Hermite polynomials in asymptotic representations of generalized Bernoulli, Euler, Bessel and Buchholz polynomials, (preprint).

[5] G. G. Lorentz, Bernstein polynomials, printed in the United States of America, 1986.

[6] L.-C. Jang, W.-J. Kim, Y. Simsek, A study on the p-adic integral representation on $\mathbb{Z}_p$ associated with Bernstein and Bernoulli polynomials, Advances in Difference Equations, 2010(2010), Article ID 163217, 6 pages.

[7] Y. Simsek, M. Acikgoz, New generating function of q-Bernstein-type polynomials and their interpolation function, Abstract and Applied Analysis, 2010(2010), Article ID 769095, 12 pages.

[8] S. Bernstein, Dėmonstration du thėorėme de Weierstrass, fondeė sur le Calcul des probabilities, Commun. Soc. Math, Kharkow, 2(1912-13), no. 13, pp 1-2.

[9] T. Kim, Barnes type multiple q-zeta function and q-Euler polynomials, Journal of physics A: Math. Theor., 43(2010) 255201, 11pp.

[10] T. Kim, Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$, Russian Journal of Mathematical physics, 16(2009), 484-491.

[11] S.-H. Rim, J.-H. Jin, E.-J. Moon, S.-J. Lee, On multiple interpolation functions of the q-Genocchi polynomials, J. Inequal. Appl. 2010(2010), Article ID 351419, 13 pages.